




# Silk Friends

The silkworm, the mulberry tree, and instructions for the educational rearing kit





**AGREEMENT NO 101095188** 



AND EUROPEAN SCALE

# The ARACNE project

The ARACNE project (an acronym for 'Advocating the Role of Silk Art and Cultural Heritage at National and European scale'), is named after the weaver transformed into a spider by the goddess Athena, in Greek mythology, and makes silk the common element of pan-European culture and history.

ARACNE aims to contribute to the creation of a broad and interconnected ecosystem of innovation related to silk in Europe, as an instrument of expression of cultural heritage and landscape, combining culture, tradition and new industrial production in an ideal network of exchanges and visions.

ARACNE started in 2023 and has a duration of 36 months, involves 11 partners and 3 ASSOCIATED PARTNERS from 7 European and non-European countries. With a budget of around EUR 3 millions, the project aims to achieve the following specific objectives:

## Objectives:

- Enhancement of knowledge and memory for the renaissance of a European Silk Innovation Ecosystem
- Co-creation of
  human-centred and
  place specific
  creative silk-based
  solutions leveraging
  on digital
  technologies
- Implementation of innovative strategies, governance and financing models for the involved organisations
- Support the
  establishment of a
  European Silk Route,
  based on the
  tangible and
  intangible silk
  cultural heritage
- Raise awareness of
  ARACNE results,
  impacts and
  expectation for the
  constitution of a
  European Silk Route
- Enhance the
  European cultural
  identity and
  strengthen
  European
  competitiveness for
  a resilient society
- Contribution to the
  European Green
  Deal, the New
  European Bauhaus
  and the Sustainable
  Development Goals

# The Silk Friends

# **Educational paths**

The European project ARACNE has among its objectives the involvement of students in the research and dissemination of silk-related culture.

To this end, several educational different paths have been proposed. For primary and secondary schools an EDUCATIONAL KIT has been developed that allows the rearing of silkworms in the classroom on an artificial diet to keep the larvae under controlled temperature and humidity. The life cycle of the insect can also be carried irrispectively of the usual rearing season (springtime), at times when there are no leaves on the mulberry tree, the only food of the silkworm. The kit is reusable and allows students to understand the process of insect metamorphosis and to become directly familiar with the developmental stages of the silkworm.



# **Educational aims**

- Skills of observation
- Development of scientific curiosity
- Show how nature and human activity are connected
- Encourage empathy towards living beings

- encourage children to closely study the development of the silkworm and to identify the mulberry plant.
- stimulate interest in the natural sciences through exploring and sensory experiences.
- teach children about the historical and economic role of the silkworm and how it is linked to the culture and history of their region.
- help children to understand how to have a kind and respectful relationship with animals and plants.

# Table of Contents

	Introduction	03
	→ The ARACNE project	03
	Educational paths	04
	Educational aims	04
01	An amazing insect	07
	1.1 Physiological characteristics of <i>Bombyx mori</i>	08
	1.2 Larva	09
	1.3 Chrysalis and cocoon	11
	1.4 Moth	13
	1.5 Life cycle	14
02	1The educational kit: rearing in the classroom	15
	2.1 What the KIT contains	16
	2.2 Preparation: what you need	17
	2.3 Good practices: observing and monitoring	18
	2.4 Egg incubation	20
	2.5 Larval hatching	21
	2.6 Feeding larvae	22
	2.7 Moulting	25
	2.8 Spinning phase	28
	2.9 Cocoons and moths	32
	2.10 Kit cleaning	35

03	The silkworm origin	37
U U	3.2 Silk discovery	38
	3.3 Legends from around the world	40
04	The Silk Road	43
UT	4.1 The long journey from China to Europe	44
	4.2 Marco Polo and the Silk Road	45
05	Mulberry	47
UJ	5.1 <i>Morus nigra</i> and <i>Morus alba</i>	49
	5.2 The silkworm's only food	50
$\cap$	The economics of the silkworm	51
06	6.1 Silk	52
	6.2 Silk processing	53
	6.3 Alternative uses of silk	54
<b>07</b>	Educational Workshops	55
UI	7.1 Suggestions for educational workshops	56
	Bibliography	61
	Credits	62





# O1 An amazing insect



# 1.1 Physiological characteristics of Bombyx mori

The silkworm (*Bombyx mori*) is an insect of the order Lepidoptera and the family Bombycidae.

Scientific classification:

Kingdom: Animalia Phylum: Arthropoda

• Class: Insecta

Order: LepidopteraFamily: BombycidaeGenus: Bombyx

• Species: mori

It is a **holometabolous** insect, which means that it undergoes complete metamorphosis. This means that it goes through four different stages in its life cycle, during which it completely changes its appearance and anatomy.

## Egg from which the silkworm hatches.

Larva (worm)
the actual silkworm,
which grows and
feeds on mulberry
leaves.


Chrysalis or Pupa the intermediate stage of transformation in the cocoon. Adult (moth) which is the mature insect ready to reproduce.












# 1.2 Larva

Larvae hatch from eggs about 1 mm in diameter, the colour of which is usually grey but can vary among strains. When they are laid, they are mainly yellow, then turn brown, then grey in about 72 hours. At the end of the incubation period, the egg whitens (becomes lighter) because the embryo, which is about to hatch, feeds on the dark membrane, which surrounds it, and protects it (in nature) from the sun's rays.

At birth, the larva is about 2 mm long, dark and hairy.



#### Aracne project Agreement NO 101095188

Like other insects, the body of the larva consists of three parts: head, thorax and abdomen. The head is dark and contains the mouthparts, which allow the silkworm to feed on mulberry leaves. The thorax has a small protuberance, and many strains have dark spots resembling a mask or eyes ("eye spot"). There are three pairs of legs on the underside, which the larva mainly uses to bring the leaf to its mouth. The abdomen takes up most of the body: on the upper part we can see the segments and dark star spots, while on the lower part there are five pairs of prolegs, which allow the larva to climb. All along the body we find the spiracles, which are small openings that allow the larva to breathe.





The mature larva measures about 8-9 cm and increases its weight 8000 times since hatching. As the silkworm prepares to spin, i.e. to find an ideal place to build its cocoon, it turns yellowish and becomes slightly transparent as it stops absorbing nutrients from the leaves and expels the remains of its last meal. The silk is produced by two glands on either side of the body and excreted through an opening under the mouthparts called spinneret.




# 1.3 Chrysalis and cocoon

Cocoon is completed in two to three days and is made up of a single continuous silk thread, the length of which varies depending on the strain and can be up to 1,500 metres.

The colour also varies from strain to strain, ranging from orange to bright yellow, pink and pale green. This is because the digestive systems of the larvae of different strains allow different amounts and qualities of the pigments in the mulberry leaf to pass through to the silk glands and from there to the thread. The colour is lost in the industrial processing of cocoons, partly because it is washed away with hot water and partly because it is sensitive to light and heat.






The function of cocoon is to protect larva while it transforms into chrysalis and then into moth. This transformation is a process that makes the animal particularly vulnerable.

Before the start of egg production process, the chrysalises are removed from cocoons after a week from spinning and divided into males and females. In order to identify the sex, it is necessary to carefully observe the abdomen, which has special markings that make it possible to separate the males from the females and then to control the mating in order to plan the necessary crosses to obtain more productive breeds, such as the poly-hybrid used for commercial rearing (and also used in the educational kit), obtained by crossing 4 breeds.







# 1.4 Moth

Moth is creamy white and about 5 cm long. It has two pairs of wings that do not allow it to fly. It emerges from the cocoon thanks to a secretion that dissolves the glue holding the silk threads together, allowing it to break through.

Like many other lepidopterans, the silkworm moth cannot feed because it no longer has the organs needed to absorb nutrients. As a result, it lives only few days for the sole purpose of reproduction.

Females can be distinguished from males on the basis of the size of their abdomens, which are much larger in females, which contain the eggs.




Adults mate for few hours. At the end of the mating they separate and the female lays about 500 eggs.

# 1.5 Life cycle

The biological cycle of silkworm lasts between 45 and 50 days in total, depending on environmental conditions (temperature and humidity affect the duration of each phase).

The larval phase is the longest and lasts about 24 days in total, divided into five instars, separated by four moults.

Counting of days always refers to the first feeding after birthing or moult.




# 02

# The educational kit: rearing in the classroom





## 2.1 What the KIT contains




The DIDACTIC KIT consists of the following materials that must be partly assembled:


- Large plexiglass container with cavity (in which to carry out the rearing)
- Adapter ring for plexiglass lid
- Circular perforated plexiglass lid
- Plexiglass hole cover (to be mounted on the lid)














- Electric heater
- 1 container with silkworm eggs
- 15 containers with diet





The material is reusable after being washed (see chapt. 2.10)



# 2.2 Preparation: what you need

To conduct silkworm rearing, it is necessary to obtain the following materials:

- Plastic cutlery or spatula (for portioning the diet)
- 2 Small transparent plastic containers with lids (approx.
   10x15 cm)
- Bottle stopper
- Oven paper
- Blotting paper (roll)
- Magnifying glass
- Straw paper grammage 100 (alternatively non-glossy parcel paper or opaque white cardboard)
- Cardboard cutouts
- Phillips screwdriver (for lid assembly)
- Microfibre cloth or non-abrasive sponge (for cleaning the kit)
- IPA cleaner Isopropyl alcohol (for kit cleaning no ethyl or 'pink' alcohol) or alcohol-free glass cleaner
- Disposable gloves\*
- Precision balance (at least one decimal)\*







\*Not mandatory

# 2.3 Good practices: Observing and monitoring

Few simple rules to follow for successful rearing:



 CLEANLINESS: always wash your hands thoroughly and sanitise tools after use.
 A clean environment minimises the risk of the silkworm getting sick.



 OBSERVATION: as the silkworm is a living being, it is necessary to observe its behaviour and appearance. The indications on the timing of the various stages are subjected to variations that can be caused by many factors (changes in temperature, excessive humidity, diseases). It is important to look closely at the eggs and larvae with a magnifying glass to identify the time of hatching and of moulting.



• MONITORING: check the growth of larvae on a daily basis and record your observations. Control temperature and humidity. To monitor larvae development is important to count the days from the administration of the first meal.

**NOTE:** Identify the correct period to start rearing. The activity cannot be carried out during periods of long breaks for holidays and long weekends, as it requires constant supervision. The kit can be left unattended for a maximum of 48 hours, taking care to check and possibly administer the meal for both days (weekend). If it is not possible to be present at all times, we recommend that you take the kit home with you for as long as necessary, taking great care during transport.



# 2.3.1 Prepare the kit to accommodate the eggs

- Remove the protective film from the plexiglass parts of the kit and clean them (see Chapter 2.10 *Cleaning the kit*).

  Dry the kit with paper towels.
- Fasten the cover with the ventilation holes on the screw and the supplied nut using the star-shaped screwdriver, taking care not to overtighten to prevent rotation of the component and to avoid damaging the disc.
- Place the electric heater into the cavity of the large container by inserting the wire into the groove provided.
- Place the thermometer inside the large container. Close with the cover, insert the plug of the heater and adjust the thermostat until the temperature inside stabilises at 25°C.



### 2.3.2 Putting the kit into operation

- Cover the base of the small transparent plastic container with a sheet of oven paper cut to size.
- Insert the bottle cap inside with a folded piece of absorbent paper soaked in water.
- Place the eggs in the small container and place its lid without closing it, leaving a gap for air to pass through.
- Place the small box in the centre of the container and close the lid, keeping the ventilation holes closed too.

# 2.4 Egg incubation

The container of the silkworm eggs is marked with the date of dispatch. The incubation period can vary considerably depending on the temperature conditions to which the eggs are exposed during their journey. Low temperatures slow down the development of the embryo (see Chapter 2.3.2 for the positioning of the eggs in the kit).

The first larvae (or young silkworms) should appear within a time window of 10 to 15 days from the date of shipment. To ensure incubation under optimum temperature conditions it is advisable to place the eggs in the kit as soon as they are received and follow the instructions for setting up the equipment and start rearing.





# 2.5 Larval hatching

When the eggs begin to hatch, you will see a few larvae, small millimetres in size, covered with dark bristles. The light-coloured shells of the eggs from which larvae have hatched are clearly distinguishable from the unhatched, dark-coloured eggs.

Hatching takes about 24 hours from the appearance of the first hatchlings. After one day, the newborn larvae should be separated from the unhatched eggs and counted. Empty shells and unhatched eggs should be disposed of in the organic waste.

After hatching, remove the cap with the wet paper from the kit.





# 2.6 Feeding the larvae

The educational kit includes jars of DIET to be fed to the worms. Each container is labelled with the instar to which the food is intended. The food should be stored in the fridge like any other perishable food.

• First instar 1 jar of diet for the entire duration

• Second instar 1 jar of diet for the entire duration

• Third instar 1 jar of diet for the entire duration

• Fourth instar 2 jars of diet, one at the beginning and one at the

middle of the development

• Fifth instar 8 small doses of diet to be given daily

Two extra diet jars are also included.

From the first to the third instar the silkworm must be reared in the small transparent plastic container with the lid.

The diet is sufficient for feeding a maximum of 20 larvae in the fifth instar. It is necessary to carry out a selection during the molting phases.



First instar: newly-hatched larvae



Third instar: larvae on the diet



Fifth instar: larvae on the diet



Second instar: larvae on the diet



Fourth instar: larvae on the diet



Fifth instar: larva during the spinning phase

#### 2.6.1 How to give the diet

The diet should be given at **room temperature**. Take the jar out of the fridge and wait half an hour.

Chop up the diet with plastic spatula and distribute it evenly on new, clean oven paper in the rearing container (small plastic container for the first instars or large tub from the kit for the last two instars). The food should be placed in small portions very close each other and not too spread\* out on the bottom of the box, or will dry out quickly.

Use the brush\*\* or plastic tweezers to transfer the worms to the new diet.

**Caution:** worms produce silk to attach themselves to surfaces. It may be useful to use a stick (such as a coffee stick) to gently separate the larvae from the food and tools.

The first meal should be given 24 hours after the first eggs have hatched. The larvae should be picked up with the brush and gently placed on the food.

Do not feed during the moult. It is important to stop feeding the larvae until the end of this physiological phase. \* From the first to the third instar, the diet shall not be too spread (take as example the image hereunder on the left), while in the other instars, it may be more spread (as in the image hereunder on the right).

\*\*The fine brush is used to shift the larvæ only in the first instar. DO NOT USE the tweezers on the newly hatched silkworms





# 2.7 Moulting

Moulting is a delicate stage in the development of larvae. Care must be taken during rearing to identify the moment when the first worms are about to moult, by detecting the signs:

- the larva stops eating
- the larva does not move and remains in a fixed position with the thorax upwards and the head downwards.

It is important **not to touch the moulting worm** in order not to compromise the process. To free itself from the exuvia (the old skin), the larva anchors itself to the surface. There is a risk, if you touch it, to damage the anchorage and compromising escape from the exuvia.

When the first worms have completed their moult you can separate them from those still moulting.

To distinguish a moulting worm from a still-moulting or not moulted worm, you must look carefully at the head (cephalic capsule).



Moulting worms have much larger heads (blue circle), almost twice the size of the non-moulted worms (yellow circle).

Use tweezers to move the moulted larvae and gently, with the help of the stick, place them into a separate plastic box without food and remove it from the kit.

Keep the moulted worms fasting for 24 hours at a temperature of not less than 20 °C.

This step allows rearing the larvae in a synchronised manner without disrupting their growth, as unfed larvae at low temperatures 'suspend' their development temporarily.

The moulting larva, on the other hand, is characterised by being motionless, not feeding, and by the presence of a triangle on its head, similar to a hood, which is formed by the capsule that is about to emerge and pushes the old one forward until it detaches (in the picture circled in green). The new head capsule is clear because the chitin that forms the exoskeleton is still soft and gradually becomes darker while hardening.

NOTE: If in doubt, it is preferable not to touch the larvae and only move them when the end of the moult has been determined.







After 24 hours, use tweezers to remove the remaining moulted worms and transfer them into the box without the food, along with the others.

Discard the old food and the worms that have not moulted, as they may be diseased and may not survive\*.

Clean the tray, place a new sheet of greaseproof paper and distribute the new food.

Use tweezers to move the moulted larvae onto the new food and put everything back in the kit at a temperature of 25  $^{\circ}$ C.

The daily count can be resumed from the first feeding. The larvae will then all be at the same growth stage.

**Caution**: The moult between IV instar and V instar takes 48 hours. Continue to separate the moulted worms on both days.

# 2.8 Spinning phase

At the end of the fifth instar, the larva has reached maturity: it measures 8-9 cm and has increased its weight 8000 times since hatching. Once they have reached full development, the worms prepare to spin.

This stage can be identified by the following signs:

- The larva stops feeding
- The larva becomes paler, yellowish and slightly transparent.
- The larva purges (gut emptying) so that the excrements appear green and softer, as the worm stops absorbing nutrients (while the black and solid ones are produced during the growth phase).
- The larva tends to climb upwards in search of a place where to build the cocoon.




In order to spin its cocoon, the silkworm must construct a structure that is anchored to 3 support points at the right distance from each other. Its instinct leads it to find the ideal place upwards, away from the dampness of the bottom of the box.

Therefore, before placing the spinning frames in the box, it is necessary to prepare the bed, i.e. the bottom of the box. Remove the food and clean the kit with paper towels, trying to remove faeces and food remains. Place a piece of cardboard on the bottom, some straw paper (or opaque wrapping paper) and a few sheets of blotting paper so that the entire bottom of the kit is covered. It is necessary to line the bottom of the kit to absorb excess moisture and prevent the formation of mould




### There are different ways to build the spinning frames:





The spinning frames in commercial rearing for cocoon production:







# 2.9 Cocoons and moths

When the larvae climb to the frames, they start spinning their own cocoons. It takes about 2-3 days to complete them. After a week, it is possible to move the cocoons, as the transformation of the larva into a pupa has already occurred, and there is no risk of compromising the development of this process.

Open the kit and gently remove the spinning frames, being careful not to damage any cocoons built in contact with the walls or the lid.

Carefully detach the cocoons from the spinning frame.

Clean the cocoons from the floss, which is the first strand of silk emitted by the silkworm to prepare the structure on which to spin the cocoon.

This silk floss can also be used in certain production sectors, for example to create padding and to extract silk proteins for the cosmetic or biomedical industries.





Clean the kit well with absorbent paper and a damp cloth, making sure to dry the inside of the kit completely. Line the bottom with a folded sheet of kraft paper. Cover the base and the walls of the kit with several layers of absorbent paper up to halfway, in order to prepare the place to host the moths, which are skilled climbers.

Position the thermometer again and place the cocoons inside the kit. Close the lid while keeping the ventilation holes open.



The emergence of the adult (a term that indicates the moth's exit from the cocoon) begins about two weeks after spinning.

The adult secretes a drop of liquid through its mouth opening, dissolving the glue that holds the thread folded in the cocoon, and it creates an opening with its legs among the silk fibres.

Once emerged, the moth can expel a yellowish liquid, a residue from the transformation of its internal organs. This is why it is important to properly line the kit's container with absorbent paper.





Silk moths lack a digestive system, so they cannot feed. They live for a few days solely to reproduce and ensure the continuity of the species. Male moths immediately begin searching for a female to mate with.

At the end of the mating, the female lays eggs that stick to the surface thanks to a glue substance produced by the moths itself. This can also happen on the walls of the kit.

# 6.10 Kit cleaning

Once the rearing is completed, it is time to clean the kit and dispose of the organic waste in the compost bin.

DO NOT keep the eggs\*, as without analysing the moths under a microscope, it is not possible to rule out contamination from diseases transmissible to the next generation; furthermore, without the correct preservation procedure, it would still be very unlikely for the larvae to hatch uniformly.

- Remove the heater and store it away.
- Unscrew the ventilation hole cover from the lid of the kit
- Rinse the plexiglass components with warm water and gently wipe with a non-abrasive sponge or microfiber cloth to remove any eggs or stains from the walls of the tank. A thin toothbrush can be used to assist in this process.
- Clean the internal walls of the tank with a microfiber cloth and isopropyl alcohol (or a non-alcoholic glass cleaner).
- Dry the kit thoroughly to avoid limescale deposits.
- Wash all tools with soap or detergent.

#### **CAUTION:**

DO NOT use ethanol or cleaners that contain it, as it can cloud the plexiglass, compromising the kit.

\*Eggs for a new rearing program can be required from the relevant entity for the territorial jurisdiction: in Italy, CREA-AA - the Sericulture Laboratory of Padua provides them free of charge for educational purposes. It is not recommended to obtain eggs from farms that cannot guarantee their healthiness.



# 03 The silkworm origin



#### 3.1 Silk discovery

The silkworm or *Bombyx mori* is a complete metamorphosis (holometabolous) insect belonging to the order of Lepidoptera. It does not exist in the natural state, so it depends entirely on man for its food, reproduction and survival. It is characterised by several races, distinguished on the basis of their geographical origin or by the characteristics of the larva and cocoon (colour, shape, length, life cycle, number of generations per year, silk production).

Sericulture is the rearing of silkworms by man and began with the domestication of silkworms between 7,000 and 4,000 years ago. The ancestor of the silkworm is a wild moth that still lives in Japan, Korea and China: *Bombyx mandarina*.

For thousands of years, China had guarded the secret of silk as a precious treasure, so much that exporting live worms was considered a very serious crime, punishable by death. Only around 500 A.D., thanks to a daring journey shrouded in legend, silkworm eggs were smuggled out of the Celestial Empire, bringing with them wonders and new possibilities. Since then, the silk has become a universal symbol of beauty, patience and human talent.

There are many other species of wild Lepidoptera that produce silk... Why has the mulberry silkworm spread all over the world?

Some species produce a cocoon with interrupted filaments, others produce a silk cocoon with a continuous thread, which, however, is more difficult to process and produces silk of a coarser quality in comparison to the mulberry silkworm.

Legend has it that the art of silk weaving was discovered by a Chinese empress. One day, while walking in her gardens, she noticed the cocoons that silkworms had formed on mulberry branches. She picked one up and it fell into the boiling water of her teacup. As she was observing, the cocoon began to unravel, revealing a long, shining thread. Fascinated, the empress learned to weave this extraordinary thread, which began the production of silk.

For thousands of years, China exported silk without revealing which animal it came from, so that many populations thought it came from trees.

It is said that some monks hid the eggs of the silkworm eggs in their reeds and brought them to Justinian, who spread the rearing of this insect throughout his empire.





#### 3.2 Legends from around the world

Here are some of the most important legends about the silkworm, fascinating stories handed down through the ages:

#### The princess and the secret of silk

According to a Chinese legend, the secret of silk was taken out of China by a princess married to a foreign prince. The princess, who did not want to part with the precious fibre, hid some silkworm eggs and mulberry seeds in her hair. Thanks to her gesture, sericulture spread to other regions, including Central Asia.

#### The silkworm and the moon goddess

According to some Chinese traditions, silkworms are believed to be a gift from the moon goddess Chang'e. It is said that the goddess taught humans how to rear silkworms and reel the silk to create clothes that resembled the silver light of the moon.

#### The sacrifice of the young peasant girl

An ancient Japanese legend tells of a young peasant who sacrificed her life to protect a harvest of silkworm cocoons from being attacked by evil spirits. After her death, her spirit was reincarnated as a moth, symbolising the continuity of the life cycle of the silkworm and the silk thread.



#### The magic mulberry tree

A traditional Korean fairy tale, tells the story of an old magical mulberry tree that could speak and protect silkworms. This tree, considered sacred, taught farmers how to care for the worms to produce the finest silk.

These legends emphasise the deep connection between the silkworm and human imagination, and highlight the symbolic cultural significance of this small insect.



# The Silk Road





## 4.1 The long journey from China to Europe

Silk was so beautiful and rare that the kings and emperors of countries outside China coveted it, but no one knew how it was made.

To get silk from one place to another, merchants used a very long itinerary called the Silk Road. It was not a road, but a series of long and difficult routes that crossed mountains, deserts and cities full of colour, markets and people from all over the world.

Merchants and their caravans, laden with coveted rolls of silk, were used to travel for months, selling the fabrics in exchange for gold, spices and precious stones.

Gradually, silkworm rearing spread along the Silk Road with the cultivation of mulberry trees, until, first the Arabs, and then the inhabitants of Byzantium (the capital of Justinian's Empire) expanded it to Europe. Europeans learned how to rear worms and process silk.

This fibre also began to be produced in Italy, first in the south, especially in cities like Palermo and Catanzaro, then in Lucca, Florence, Bologna and finally in Venice and Como, which became famous for their silk fabrics.




### 4.2 Marco Polo and the Silk Road

Marco Polo, a famous Venetian traveller, had a special relationship with silk. Around 700 years ago, in 1271, Marco Polo set off on an incredible journey to China, following the routes of the famous Silk Road. In fact, these routes were used to exchange silk but also spices, jewellery and many other precious goods between East and West.

When Marco Polo arrived in China, he met Emperor Kublai Khan and became fascinated by the beauty and importance of silk. He told how silk was everywhere: in the clothes of emperors, in the fabrics of curtains and even in the gifts exchanged between nobles. Returning home after years, Marco Polo brought back incredible stories about China and its fibre.

His book, The Million, was one of the first to narrate the link between East and West through the Silk Road, and helped to spread the myth of the Orient, made up of distant lands, different in customs and habits, as well as of wonderful threads of silk.







#### 5.1 Mulberry

There are mainly two species of mulberry in Europe: *Morus nigra* (the black mulberry) and *Morus alba* (the white mulberry). This plant belongs to the Moraceae family.

Other species of mulberry, such as the red mulberry (*Morus rubra*), which is native to North America, are less common in Europe.

The mulberry produces a sorosis, which is an infructescence consisting of many small fruits growing together. Each of these arises from a different flower and develops from a part of the flower called the ovary.

They grow close together on a peduncle, forming what we commonly call a mulberry fruit.

#### 5.2 Morus nigra and Morus alba

The **black mulberry** originated in the Middle East and spread to Greece and southern Italy very soon afterwards, where the warmer climate is more favourable. It was known to the Greeks and Romans, who appreciated its dark red fruits for their taste and nutraceutical benefits. The word 'nutraceutical' is a combination of 'nutrition' and 'pharmaceutical', as this discipline studies all substances in foods that can promote health, prevent disease or cure it.




The white mulberry, on the other hand, was introduced later, around the year 1400 to italy, when it was brought from China. It was primarily cultivated for raising silkworms. It has become very common throughout Europe, particularly in southern regions such as Spain, France and Italy, as it thrives in colder climates in comparison to black mulberry. Its fruit changes colour, ranging from white to pinkish and red, but it is less valuable than that of the black mulberry.

#### 5.3 The silkworm's only food

In the past, the mulberry tree was cultivated almost exclusively to feed silkworms. Today, however, it is considered useful in many ways: it is aesthetically attractive in fields and gardens, its fruit is nutritious and tasty, proteins and plant fibres can be extracted from its leaves, and some of its compounds are used in natural medicines and cosmetics. Additionally, bioplastics, which are more sustainable than traditional plastics, can be produced from the mulberry tree.

As mentioned above, silkworms feed exclusively on mulberry leaves, and during their life cycle, which lasts around a month, a single larva can consume up to 20 grams of fresh leaves, 88 per cent of which are ingested in the fifth instar. This rich nourishment enables the silkworm to grow and store what it needs to produce silk and build the cocoon that will protect it during its transformation first into a chrysalis and then into a moth.



# O6 The economics of the silkworm





#### 6.1 Silk

Silk is an animal fibre produced by two specialised glands in the body of the silkworm larva. It is composed of two main proteins: fibroin and sericin.

**Fibroin** accounts for around 75% of the thread and is insoluble in water. It is characterised by its lustre, elasticity, and tensile strength.

**Sericin**, which constitutes around 25% of native silk, surrounds the fibroin like a sheath. This acts as a kind of glue, ensuring the threads can join together to form the layers of the cocoon.

#### 6.2 Silk processing

In order to be used, silk must be unwound from the cocoon in which it is enclosed, a process known as reeling. The cocoon is immersed in hot water, which dissolves the sericin, allowing the long fibroin thread to unravel. This process takes place in factories known as 'reeling plants', which are a testament to the early industrialisation of southern European territories, the traces of which can still be seen today in the large complexes with tall chimneys that dot the agricultural landscapes, especially in northern Italy.

The thread is wound onto reels, forming skeins of raw silk. However, the skein of silk still has to be 'degummed' to remove all the residual sericin and acquire its characteristic softness.

Only the finest, whitest and most perfect cocoons can be used in the reeling process. Cocoons with small spots or imperfections can be used in the carding process, which is similar to that used for wool. This produces a silk hank, which can then be processed into a discontinuous silk thread that is less valuable than unravelled silk.







#### 6.3 Alternative uses of silk

Due to its unique properties, silk has a wide range of applications beyond textiles. As a protein fibre, it can be dissolved and regenerated for use in biomedicine or as a biomaterial. In fact, silk is biocompatible, meaning it can be used to repair or replace tissues within the human body.

It is used to make membranes, tubes and nets that can be implanted in the body, for example for vascular grafts or as a coating for implants.

Fibroin can be used to create scaffolds on which human cells can grow, for the reconstruction of skin, bones, cartilages, nerves or blood vessels.

Sericin is used in cosmetics, such as creams, shampoos, conditioners, serums and ointments.

The chrysalis contained within the cocoon can be used as feed, but under certain conditions it can also be used as food for humans due to its richness in nutrients. Each byproduct can therefore become the raw material for a new supply chain.



# 07 Educational Workshops



## 7.1 Suggestions for educational workshops

This final section of the rearing guide contains suggestions for manual activities to be carry out with students. The aim is to reinforce the knowledge acquired when using the kit.

These suggestions are designed to develop pupils' skills in observation and analysis, encouraging children to understand the transformation of the silkworm in detail and identify the mulberry plant.

The activities are intended to foster empathy towards living beings and encourage respect for animals and plants. Manual and hands-on activities also promote manual dexterity and creativity.

#### Mayu no Hana

Proposing the ancient Japanese art of creating flowers from silk cocoons, called Mayu no Hana (or Hanamayu), is an original and inspiring idea to engage students in a creative manual workshop in the classroom. In Japan this tradition dates back several centuries, when silkworm farmers' families were used to employ leftover cocoons to make elegant decorations and small flowers donated as gifts or sold at markets. Today, turning the cocoons into colourful flowers can be a useful teaching project for developing manual dexterity, as well as patience and attention to detail. In addition, this activity also encourages reflection on the reuse of materials. Each pupil can choose a flower, cut the petals from the cocoons, mount them on a wire, and customise their work with colours, as well as paper or fabric leaves. The finished flowers can become original gifts for families or decorations for the classroom. A Mayu no Hana workshop keeps the tradition of craftsmanship to life, stimulating creativity while emphasising the importance of manual work in a technology-dominated age.



#### Creating animals and other decorations with silk cocoons

In addition to creating flowers, cutting out the cocoons allows you to make an infinite variety of small artefacts, such as cute little animals. You can also make wearable accessories such as necklaces and hairpins, as well as home decorations and greeting cards. With fabric or vinyl glue, the possibilities are endless.





#### Mulberry leaf herbarium

To preserve mulberry leaves for a long time, they can be used to make a classroom herbarium. This stimulates observation and attention to detail, helping children recognise the plant in their environment. This workshop has scientific and historical value, but above all it has practical meaning because it helps children to become familiar with botany, rediscover their connection with nature, and develop their organisational skills.

While waiting for the leaves to dry, you can exploit the 'frottage' technique, which involves rubbing. Simply place a white sheet of paper over the leaf, rib side up, and rub the crayon over the surface to make the details appear on the paper.

#### Pre-cinema optical games

The *thaumatrope* is an ancient optical toy consisting of a disc with two different images on each side. When rotated quickly, the images of worms, cocoons and moths merge to create a single moving scene. This tool allows you to experience the persistence of the vision effect, combining science, creativity and fun. It is an interactive experience that stimulates manual dexterity and teaches while providing entertainment.





Download thaumatrope template from the QR code

#### Building the silkworm on a leaf


You can find numerous online tutorials explaining how to make a cute larva that moves along a cardboard leaf. You can make it by using a napkin or a paper handkerchief, or by cutting out a straw. The various components are easily assembled with glue.







Bibliography:	Lingua
Fila, Gianni, and Silvia Cappellozza, eds. <i>The Silkworm:</i> History, Culture, Traditions and Science. A Commented Catalogue of the Historical Silk Science and Technology Collections Housed at the Former Experiment Sericulture Station of Padua. Rubano (PD): Turato Publisher, 2023.	English
Dedos, Skarlatos. <i>Σηροτροφία (Sericulture)</i> . Kallipos University Publications, 2023.	Greek
Centro Studi Memorandum. <i>Il bravo allevatore: Manuale.</i> Cortale: Centro Studi Memorandum, 2017.	Italian
Cappellozza, Silvia. <i>Manuale di buona pratica agricola per la bachicoltura</i> . Published online on masaf.gov.it, 2010.	Italian
Moraschini, Paolo, Lorena Cirnigliaro, and Laura Ghidini. "L'allevamento dei bachi da seta: un'attività sperimentale alla scuola primaria." Emmeciquadro, no. 29 (April): Euresis, 2007.	Italian
Grekov, Dimiter, Euripidis Kipriotis, and Panomir Tzenov.  Sericulture Training Manual. Komotini: Agricultural Research  Station of Komotini, NAGREF, 2005.	English
Faragò, Silvio, and Walter Filz. <i>Il baco da seta.</i> Milano: Stazione Sperimentale per la Seta, 2001.	Italian
Lim, Soo-Ho, Young-Taek Kim, Sang-Poong Lee, In-Jun Rhee, Jung-Sung Lim, and Byung-Ho Lim. <i>Sericulture Training Manual</i> . FAO Agricultural Services Bulletin 80. Rome: Food and Agriculture Organization of the United Nations, 1990.	English



#### **Credits:**

Text: Silvia Cappellozza, Diana Mantegazza, Graziella Paglia

and Alessio Saviane

Graphics: Diana Mantegazza

Illustrations: @irasutoya via Canva and Diana Mantegazza Photos: CREA-AA; PIOP; Paolo Zerbato from Pixabay; ivabalk from Pixabay; LoggaWiggler from Pixabay; Rafael Ben-Ari

from AdobeStock

More info: aracneproject.eu

Contact: education@aracneproject.eu

Padova - 2025





Scan for the full guide available in multiple languages https://aracneproject.eu/educational-kit/



